

MULTIWAVELENGTH OBSERVATIONS OF GAMMA-RAY BURSTS: TOWARDS THE UNDERSTANDING OF THE MYSTERY

Alberto J. Castro-Tirado

Laboratorio de Astrofísica Espacial
y Física Fundamental (LAEFF-INTA)
P.O. Box 50727, E-28080, Madrid, Spain
e-mail address: ajct@laeff.esa.es

ABSTRACT

Since their discovery in 1973, Gamma-Ray Bursts (GRBs) have remained for many years one of the most elusive mysteries in High Energy-Astrophysics. The main problem regarding the nature of GRBs has usually been the lack of knowledge of their distance scale. About 300 GRBs are detected annually by BATSE in the full sky, but only a few of them can be localized accurately to less than half a degree. For many years, follow-up observations by other satellites and ground-based telescopes were conducted, but no counterparts at other wavelengths were found. The breakthrough took place in 1997, thanks to the observation by *BeppoSAX* and *RossiXTE* of the fading X-ray emission that follows the more energetic gamma-ray photons once the GRB event has ended. This emission (the afterglow) extends at longer wavelengths, and the good accuracy in the position determination by *BeppoSAX* has led to the discovery of the first optical counterparts -for GRB 970228, GRB 970508, and GRB 971214-, greatly improving our understanding of these puzzling sources. Now it is widely accepted that most bursts originate at cosmological distances but the final solution of the GRB problem is still far away.

Key words: multiwavelength observations; gamma-ray bursts.

1. INTRODUCTION

In 1967-73, the four *VELA* spacecraft (named after the spanish verb *velar*, to keep watch), that where originally designed for verifying whether the former Soviet Union abided by the Limited Nuclear Test Ban Treaty of 1963, observed 16 peculiarly strong events (Klebesadel, Olson and Strong 1973, Bonnell and Klebesadel 1996). On the basis of arrival time differences, it was determined that they were related neither to the Earth nor to the Sun, but they were of cosmic origin. Therefore they were named cosmic γ -ray Bursts (GRBs hereafter).

GRBs appear as brief flashes of cosmic high energy

photons, emitting the bulk of their energy above ≈ 0.1 MeV (Fig. 1). They are detected by instruments somewhat similar to those used by the particle physicists at their laboratories. The difference is that GRB detectors have to be placed onboard balloons, rockets or satellites. In spite of the abundance of new observations of GRBs, their energy source and emission mechanism remain highly speculative.

The KONUS experiment on *Veneras 11* and *12* gave the first indication that GRB sources were isotropically distributed in the sky (Mazets et al. 1981, Atteia et al. 1987). Based on a much larger sample, this result was nicely confirmed by BATSE on board the *CGRO* (Meegan et al. 1992). About 300 GRBs occur annually in the full sky, but only few of them are localized accurately. The apparent isotropy was interpreted in terms of GRBs produced at cosmological distances, although the possibility of a small fraction of the sources lying nearby, within a galactic disc scale of few hundred pc, or in the halo of the Galaxy, could not be discarded. Another result was that the time profiles of the bursts are very different, with some GRBs lasting a few ms and others lasting for several minutes. In general, there was no evidence of periodicity in the time histories of GRBs. However there was indication of a bimodal distribution of burst durations, with $\sim 25\%$ of bursts having durations around 0.2 s and $\sim 75\%$ with durations around 30 s. A extensive review of the observational characteristics can be found in Fishman and Meegan (1995).

A deficiency of weak events was also noticed, and all these observational data led many researchers to believe that GRBs are indeed at cosmological distances. In this case, the released energies could be as high as 10^{53} erg and models could involve coalescence of neutron stars in double systems, neutron star-black hole systems, accretion induced collapse in white dwarfs or "failed" Type I supernovae. It was also proposed the possibility of having a mixture of two populations: a cosmological plus a galactic one, but the latter, probably formed by accreting neutron stars, will account for only a very small fraction of the total population. See Nemiroff (1984) for a review of the different theoretical models.

It is well known that an important clue for resolving

the GRB puzzle is the detection of transient emission -at longer wavelengths- associated with the bursts. Here I review all the efforts in the search for GRB counterparts throughout the electromagnetic spectrum. I will first review the searches prior to 1997, and afterwards I will discuss the important discoveries achieved last year. Previous reviews can be seen in Schaefer (1994), Hartmann (1995), Vrba (1996), Greiner (1996a) and Hurley (1998).




Figure 1. One of the GRBs detected by BATSE, lasting for about 150-s. Further details in Connaughton et al. (1997).

2. SEARCHES PRIOR TO 1997

2.1. X-ray searches

2.1.1. Searches for simultaneous and near-simultaneous X-ray emission

The observation of X-rays below 20 keV is important in order to understand the nature of the physical processes in the GRB sources. Most of the experiments devoted to GRB study had lower cut-off energies of the order of 20-40 keV. However, a few of them were designed in order to cover part of the soft X-ray range.

Already in 1972, the Apollo 16 spectrometer detected X-ray photons in the 2-8 keV range for GRB 720427 (Trombka et al. 1974). Wheaton et al. (1973) observed in GRB 720514 the presence of an X-ray signal which lasted *longer* than the gamma-ray event. In 1984, Katoh et al. reported an X-ray tail for GRB 811016 that lasted more than 30 s after the termination of the hard event. *GINGA* also detected emission down to 1.5 keV, with soft X-ray tails found for 8 events (Murakami et al. 1991, Yoshida and Murakami 1994). Moreover, a significant fraction ($\sim 10\%$) of the GRBs detected by WATCH on *GRANAT* displayed X-ray tails in the 6-20 keV range (Castro-Tirado et al. 1994a). Connors and McConnell (1996) have discussed the proposal that the X-ray transient detected by *HEAO 1* following GRB 780506 could indeed be related to the burst itself. Therefore it seemed clear that, at least for some bursts, there was

evidence for X-ray emission lasting longer than the GRB.

2.1.2. Deep searches

Grindlay et al. (1982) and Pizzichini et al. 1986 initiated the searches for X-ray quiescent counterparts by means of *EINSTEIN*. Five GRB error boxes were observed and, with one possible exception (GRB 781119, Grindlay et al. 1982), no point source was detected in any of the observations. Assuming that the bursts originated from galactic neutron stars, only upper limits to the surface temperatures were derived (Pizzichini et al. 1986). Boer et al. (1988) observed the GRB 781119 region, but failed to detect the source with *EXOSAT*. Recently, *ASCA* and *ROSAT* "rediscovered" the X-ray source in GRB 781119, with a flux of $F_{(0.5-8 \text{ keV})} \sim 1.6 \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$ (Hurley et al. 1996, Boer et al. 1997). Further evidence for an X-ray source associated with a GRB was obtained by *ASCA* in the GRB 920501 error box (Murakami et al. 1996). The X-ray spectrum is heavily absorbed at low energies, implying an absorption consistent with the galactic value in that direction, and suggesting that the X-ray source is *quite* distant. Using the *ROSAT* all-sky-survey data, Greiner et al. (1995) detected 27 X-ray sources in 45 small GRB error boxes. On the basis of the optical identification of these sources, no convincing association to GRBs was found. Moreover, the number of sources corresponded to the expected number of background sources. The upper limit to the X-ray flux was $F_{(0.1-2.4 \text{ keV})} \sim 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$. Another field (GRB 940301) was observed by *ROSAT* four weeks later and revealed no fading counterpart (Greiner et al. 1996b).

2.2. EUV searches

2.2.1. Near simultaneous searches

Searches at EUV wavelengths have only been performed since 1992, following the launching of *EUVE*. Castro-Tirado et al. (1998a) has reported a serendipitous observation in the extreme ultraviolet window by *EUVE*, ~ 11 hours after the occurrence of GRB 921013b. No source was detected in the EUV range.

2.2.2. Deep searches

A deep observation on the GRB 920325 error box was performed 17 months after by *EUVE*. No quiescent source was found, thus constraining a neutron star as a possible counterpart (Hurley et al. 1995).

2.3. Optical searches

2.3.1. Archival searches

The strategy behind this kind of search is the assumption that GRBs repeat. Some scientists have assumed

that GRBs could be a repetitive phenomenon, and they have used all the available information at the major plate archives in order to look for optical transient (OT) emission in some of the smaller GRB error boxes. Atteia et al. (1985) searched for about 1500 hours but no such transients were seen. Schaefer discovered in 1981 the first OT in a Harvard plate taken in 1928 (OT 1928), located inside the small GRB 781119 error box. Another OT was found in the GRB 781006b error box in three different plates taken in 1966. A large amplitude flaring dMe star, probably unrelated to the burst, was proposed to be related to the OT (Greiner and Motch 1995). Another good candidate is OT 1905, found by Hudec et al. (1994) slightly outside the interplanetary network error box for GRB 910219. Although these OTs seem to be real, another ~ 40 candidates were found, but most of them were rejected as they turned out to be plate defects. See a discussion in Hudec (1993, 1995). However, there is no convincing proof that the OTs found in plate archives and GRBs are related to the same physical phenomenon.

2.3.2. Deep searches

Deep searches were carried out in the 80's for the smallest GRB error boxes provided by the Interplanetary Network: GRB 781119 (studied by Pedersen et al. 1983) and GRB 790406 (Chevalier et al. 1981). Recently, with the advent of the newer big CCD detectors, deep searches have been carried out of GRB 790329 and other error boxes (Vrba, Hartmann and Jennings 1995). The most powerful telescopes, such as the 6-m in the Caucasus and the Hubble Space Telescope (HST) have also been used for observing the error boxes of GRBs: GRB 790613 (Sokolov et al. 1996) and GRB 790325 and 920406 (Schaefer et al. 1997a). In any case, no quiescent GRB counterpart has been firmly established. In general, the small GRB error regions did not contain galaxies to fairly deep limits (the so-called "no-host problem", Fenimore et al. 1993), and it was obvious that the bursts, if at cosmological distances, could not come from normal galaxies. Either they had to occur in subluminal galaxies or their progenitors somehow had to be ejected from the parent galaxy.

2.3.3. Near-simultaneous searches

The first search for optical emission arising *simultaneously* with a GRB was performed by Grindlay, Wright and McCrowsky (1974). Hudec et al. (1987) and Hudec (1993) found five events for which time-correlated plates were available. Greiner et al. (1996c) found plates within ± 12 -h of the bursts, but none revealed interesting brightenings in the error box. For GRB 920824, a rather deep plate was taken simultaneously with the burst in Dushanbe. They conclude that the optical emission of typical GRBs is at level below $F_{opt}/F_{\gamma} \sim 2$ during the burst and $F_{opt}/F_{\gamma} \sim 20$ -400 a few hours after the burst, with F_{γ} and F_{opt} being the peak gamma-ray (and optical) fluxes. Stronger upper limits have been obtained by GROCE (Lee et al. 1996) who have recorded sky images for more than 30 GRB triggers, in many cases

while the burst was still in progress. Additional results were obtained by Castro-Tirado et al. (1994b) regarding follow-up observations of eight bursts detected by WATCH. No transient optical emission was detected in the following hours down to 18th magnitude. Similar observations were reported for GRB 930131 by Schaefer et al. (1994) and McNamara and Harrison (1994). Again, no firm counterpart was detected in their observations. Barthelmy et al. (1994) obtained upper limits for any transient optical emission arising from three bursts, between 35 hr and 8 days following the burst. The only OT found nearly-simultaneously to a GRB event, is the one reported by Borovička et al. (1992). They found a star-like spot at the edge of the GRB 790929 error box on a plate taken only 7.1 hr after the high energy event. Also, two additional objects were found at the same position on another plates. The objects are consistent with the star HDE 249119. Further optical studies detected additional optical eruptions of HDE 249119, reporting ten optical brightenings with amplitudes greater than 0.5 mag (Štěpán & Hudec 1996), but it is unlikely that this star would be related to GRB 790929.

2.4. Near-IR searches

2.4.1. Deep searches

The first search for counterparts in the near-IR was initiated by Schaefer et al. (1987), who investigated seven GRB error boxes at a wavelength of $2.2 \mu\text{m}$ (K-band). No convincing candidates were found, with a detection limit of $K = 19$ for two error boxes. An imaging survey of the six smallest GRB error boxes from the third Interplanetary Network has revealed relatively bright galaxies in each of them (Larson, McLean and Becklin, 1996). Although the statistical significance of their result is not overwhelming, it leaves the "no-host problem" as an open question. Klose et al. (1996) obtained deep imaging of the GRB 790418 error box, showing several galaxies as well. Deep infrared imaging of the quiescent X-ray source located within the GRB 920501 error box has been performed by Blaes et al. (1997). One of the objects is a type 1 Seyfert galaxy, and Drinkwater et al. (1997) has identified the galaxy with the X-ray source seen by *EINSTEIN*, *ASCA* and *ROSAT*, strengthening the above-mentioned results. Is this Seyfert galaxy the true GRB counterpart?

2.5. Mid-IR searches

Searches for *quiescent* counterparts have also been conducted in the past in the IR. Schaefer et al. (1987) used the IRAS data base at wavelengths of 12, 25, 60 and $100 \mu\text{m}$ and looked for candidates within 23 well localized GRB error boxes. No convincing counterparts were found.

2.6. Millimetre searches

Two simultaneous observations were obtained by the COBE satellite, when correlating GRBs that o-

occurred during an 8 month overlap period (Bontekoe et al. 1995, Stacy et al. 1996). The 2σ upper limit quoted is 31000 Jy at 53 GHz. A likelihood analysis was performed by Ali et al. (1997) in order to look for a change in the level of millimetric emission from the location of 81 GRBs during the first two years of the mission (1990-91). No positive detection was achieved, with 95 % confidence level upper limits of 175, 192 and 645 Jy respectively at 31, 53 and 90 GHz. Another team at the Fallbrook Low-Frequency Immediate Response Telescope (FLIRT) searched for prompt radio emission at 74 MHz arising from GRB sources. The delays were of the order of 2 to 12 min. Preliminary results for 12 GRBs indicated that no source was found above a ~ 100 Jy detection limit (Balsano et al. 1996).

2.7. Radio searches

2.7.1. Near simultaneous searches

A serendipitous observation towards the error box of GRB 920711 was performed at 151 MHz with the Cambridge Low-Frequency Synthesis Telescope (CLFST) from 2 to 4 days after the occurrence of GRB 920711. No source was detected down to 40 mJy (Koranyi et al. 1994). This instrument also observed GRB 940301 within 1 hour after the burst, imposing a 200 Jy upper limit, and 80 mJy between 1-36 days after the event. CLFST observations were conducted for another two bursts, with flux limits of 35 Jy for 2.5 hours after GRB 950430 and 16 Jy for 1 hour after GRB 950706 (Green et al. 1996). A search for a radio transient at 4.86 and 1.4 GHz within the GRB 930131 failed to detect any source (Schaefer et al. 1994). VLA observations were performed 23 hours after GRB 940217, but only an upper limit of $\leq 4 \mu\text{Jy}/\text{beam}$ was given by Palmer et al. (1985). No fading/flaring radio counterpart was found either at 1.4 GHz (upper limit of 3.5 mJy) or 0.4 GHz (upper limit of 55 mJy) within the GRB 940301 error box (Frail et al. 1994).

2.7.2. Deep searches

It was suggested that if GRBs were to be related to highly magnetized, rotating neutron stars, they would produce polarized radio emission during the quiescent state. In fact, VLA maps of the GRB 781119 error box showed two sources (Hjellming and Ewald 1981). One showed some degree of polarization, and the other coincided with the *ROSAT* X-ray source (Hurley et al. 1996). Schaefer et al. (1989) found four radio sources in 10 small GRB error regions. None of them displayed unusual properties and the number of sources was the expected number of chance background sources. They concluded that none of them were associated with GRBs. Deep VLA observations of three GRBs were also performed by Palmer et al. (1995). GRB 920501 was observed 13 days and 9 months after the burst, at frequencies of 1.4 and 8.4 GHz, imposing upper limits of $\leq 235 \mu\text{Jy}/\text{beam}$ and $\leq 77 \mu\text{Jy}/\text{beam}$ respectively. A strong limit was derived for GRB 930706 as $\leq 1 \mu\text{Jy}/\text{beam}$ after 9 days.

2.8. VHE/UHE searches

Photons with energies up to 18 GeV have been detected in only one case, 1.5-hr after GRB 940217 (Hurley et al. 1994). There are models that predict TeV emission from GRBs (Meszaros, Rees and Papathanassiou 1994, Halzen et al. 1991). A positive detection would constrain models with respect to the emission mechanism and the distance, because no TeV photons are expected if they occur at cosmological distances due to the interaction of the photons with the microwave and infrared background. This process leads to pair production and might become important at $E \geq 30$ TeV, as pointed out by Wdowczyk, Tkaczyk and Wolfendale (1992), Mannheim, Jartmann and Funk (1997) and Stecker and de Jager (1997). In fact, at $E \geq 50$ TeV, there seems to be evidence of a detection from the Dublin ASA (GRB 910511, Plunkett et al. 1995). The rest of the current arrays and Cerenkov telescopes have provided only upper limits at this energy threshold. A systematic search was conducted at AES-KGF (Bhat et al. 1996) for PeV γ -rays from the locations of about 39 high fluence GRBs during Apr 91-Mar 93. Upper limits of 2×10^{-12} photons $\text{cm}^{-2} \text{s}^{-1}$ were obtained. Another search was performed for 115 GRBs at the CASAMIA detector. Flux limits of 6×10^{-12} photons $\text{cm}^{-2} \text{s}^{-1}$ at $E \geq 160$ TeV were obtained (Catanese et al. 1996). Follow-up searches have also been carried out with HEGRA above 1 TeV for more than 150 bursts, but no significant excess was detected by Krawczynski et al. (1996) or Padilla et al. (1997). In the 0.25-4 TeV range, it has been suggested that the Whipple Observatory can detect, under favourable conditions, ~ 1 event per year (only ~ 0.01 per year in the case of the HEGRA array). However, only upper limits have been set up by the Whipple telescope (Connors and McConnell 1996) and the EAS-TOP ASA experiment at $E \geq 10$ TeV (Agglietta et al. 1995).

2.9. Neutrinos and muons

The search for neutrinos, antineutrinos and muons arising either from intrinsic GRB mechanisms as proposed by Mészáros and Rees (1993) and Paczynski and Xu (1994) or Earthrock neutrino-produced muons, has been performed at several places, but none of the experiments have detected neutrinos or muons in excess of the expected background in correlation with a GRB. These are the Mount Blanc LSD telescope (Agglietta et al. 1995), the Irvine-Michigan-Brookhaven (IMB) detector (LoSecco 1994, Becker-Szendy 1995) and the Soudan-2 muon experiment (De Muth et al. 1994). Upper limits are $6 \times 10^{11} \text{ cm}^{-2}$ (for ν_e, ν_μ, ν_τ at $20 \leq E_\nu \leq 100$ MeV) and $2.1 \times 10^{13} \text{ cm}^{-2}$ (for $\bar{\nu}_e, \bar{\nu}_\mu, \bar{\nu}_\tau$ at $20 \leq E_\nu \leq 100$ MeV) (Agglietta et al. 1995).

3. 1997: THE DETECTION OF THE FIRST COUNTERPARTS AT OTHER WAVELENGTHS




Figure 2. The X-ray afterglow of GRB 970228. Based on the WFC, BeppoSAX and ASCA data. Adapted from Costa et al. (1997).

3.1. The X-ray afterglow

With the launching of *BeppoSAX*, it was possible on 28 Feb 1997 to detect the first *clear* evidence of a long X-ray tail -the X-ray afterglow- following GRB 970228. See figure 2. The X-ray fluence was $\sim 40\%$ of the gamma-ray fluence, as reported by Costa et al. (1997). Therefore, the X-ray afterglow is not only the low-energy tail of the GRB, but also a significant channel of energy dissipation of the event on a completely different timescale. Another important result was the non-thermal origin of the burst radiation and of the X-ray afterglow (Frontera et al. 1998).

Further X-ray afterglows were observed by *BeppoSAX* in GRB 970402 (Piro et al. 1997a), GRB 970508 (Piro et al. 1997b), GRB 971214 (Antonelli et al. 1997) and probably in GRB 970111 (Feroci et al. 1998) and GRB 971227 (Sofitta et al. 1997). Another three were observed by *RossixTE* in GRB 970616 (Marshall et al. 1997a), GRB 970815 (Smith et al. 1997) and GRB 970828 (Marshall et al. 1997, Remillard et al. 1997). The latter was also detected by *ASCA* (Murakami et al. 1997) and *ROSAT* (Greiner et al. 1997). The X-ray spectrum as observed by *ASCA* is strongly absorbed ($N_H = 10^{22} \text{ cm}^{-2}$, Murakami et al. 1998), suggesting that the event occurred in a dense medium.

3.2. A search for EUV counterparts

Boër et al. (1997b) reported an observation by EUVE ~ 20 hr after GRB 971214. No source was detected at 100 Å. Assuming $N_H = 10^{20} \text{ cm}^{-2}$, the upper limit to the flux is $1.7 \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$.

3.3. The optical counterparts

The improvement of GRB localizations by *BeppoSAX* (in combination with the *Third Interplanetary Network*




Figure 3. The OT related to GRB 970508, as seen in this R-band image taken by C. Wolf and R. Fockenbrock with the 2.2m telescope at the German-Spanish Calar Alto Observatory on 9 May 1997. North is up and east to the left.

of satellites) has led to the detection of the first three optical counterparts in 1997. The first optical transient associated to a GRB was identified on 28 Feb 1997, 20 hr after the event, by Groot et al. (1997). The OT was seen in earlier images taken by Pedersen et al. (1997) and Guarnieri et al. (1997), in the rising phase of the light curve. The maximum was reached ~ 20 hr after the event ($V \sim 21.3$), and followed by a power-law decay as $t^{-1.2}$ (Galama et al. 1997, Masetti et al. 1998). An extended source was seen at the OT position since the very beginning by both ground-based and *HST* observations (van Paradijs et al. 1997, Sahu et al. 1997). New *HST* observations taken 6 months after the event were reported by Fruchter et al. (1997). Both the OT (at $V = 28$) and the extended source ($V = 25.6$) were seen. The extended source surrounding the point-source was interpreted as a galaxy, according to the similarities (apparent size, magnitude) with objects in the *HST* Deep Field.

The second optical transient is associated with GRB 970508 and was discovered by Bond (1997). It was observed only 3 hr after the burst in unfiltered images (Pedersen et al. 1998a) and 4-hr after in both the U and R-bands (Fig. 3). It displayed a strong UV excess (Castro-Tirado et al. 1998, Galama et al. 1998). The optical light curve reached a peak two days after ($R = 19.7$, Djorgovski et al. 1997) and was followed by a power-law decay as $t^{-1.2}$ (Fig. 4). The long-lasting OT was observed at many ground-based observatories (Chevalier et al. 1997, Garcia et al. 1997, Mignoli et al. 1997, Schaefer et al. 1997b, Sokolov et al. 1998). Optical spectroscopy was performed at Calar Alto and La Palma (Castro-Tirado et al. 1997a) and Hawaii (Metzger et al. 1997a). The latter spectrum allowed a direct determination of the GRB 970508 redshift ($z \geq 0.835$) and was the first proof that at least a fraction of the GRB sources lie at cosmological distances (Metzger et al. 1997b). Unlike GRB 970228, no host galaxy has been seen so far, but the optical spectroscopy and the flattening of the decay in late August (Pedersen et al. 1998b) suggest that the contribution of a constant bright-

ness source -the host galaxy- has been detected. The luminosity of the galaxy is well below the knee of the galaxy luminosity function, L^* , and the detection of deep Mg I absorption and strong [O II] emission - both indicative of a dense medium- suggests that the host could be a dwarf, blue, rapidly forming starburst galaxy (Pian et al. 1998).




Figure 4. The light curve of OT/GRB 970508 in the R-band. Based on observations obtained at La Palma, Calar Alto and Loiano. The horizontal dashed line indicates the magnitude of the constant source (the galaxy) that is believed to host the GRB. Adapted from Castro-Tirado et al. (1998a).

The third optical transient is related to GRB 971214 and was identified by Halpern et al. (1997) as an $I = 21.2$ object that faded 1.4 mag in 1 day. Independently, this object was also noticed as suspicious by Itoh et al. (1997). Further observations proved that the decay followed a power-law decline (Diercks et al. 1997, Castander et al. 1997) with $t^{-1.4}$, similar to, but steeper than, the other two GRBs with optical counterparts.

In spite of intensive searches, no optical counterparts were found within one day of the event for GRB 970111, GRB 970402 or GRB 970828 (Castro-Tirado et al. 1997b, Pedersen et al. 1998b, Groot et al. 1998).

3.3.1. Near-infrared counterparts

Fading near-IR counterparts were observed for GRB 970228, GRB 970508 and GRB 971214. For GRB 970228, the IR-flux declined from $J = 21.0$ ($K \geq 19.5$) on Mar 18 (Klose and Tuffs 1997) to $J = 23.5$ ($K = 22.0$) on Mar 30 (Soifer et al. 1997). In the case of GRB 970508, the IR flux decayed from $K = 18.4$ to $K = 19.2$ in three days (Morris et al. 1997). GRB 971214 was detected in the K-band 4 hr after the burst (Gorosabel et al. 1998) and decreasing from $J = 20.3$ on Dec 15 to $J = 21.5$ on Dec 16 (Tanvir et al. 1997).

3.4. A search for mid-IR counterparts

Following the detection by BeppoSAX of GRB 970402, a Target-of-Opportunity Observation by *ISO* was performed 55-hours after the event, in the ISOCAM 8-15 μm (IRAS-equivalent bandpass) and ISOPHOT 174 μm band, in order to monitor the entire BeppoSAX GRB error box. A further observation was carried out 8 days later, with a similar set-up, in the X-ray afterglow error box. About 50 sources were detected in a field centered on the GRB error box. Seven of them lie within the smaller X-ray error circle. With the exception of the variable star BL Cir, no other variable object was found in the entire GRB error box. $5\text{-}\sigma$ upper limits for any new object are $F_{(12\mu\text{m})} \leq 0.14$ mJy and $F_{(174\mu\text{m})} \leq 350$ mJy (Castro-Tirado et al. 1998b). A negative result was also obtained in the same bandpass for GRB 970508 (Hanlon et al. 1998).

3.5. Detection in the millimetre range

GRB 970508 was detected as a continuum point source at 86 GHz with the IRAM Plateau de Bure Interferometer. A ~ 2 mJy flux was observed on 19-21 May 1997. The object was not detected any more after 28 May. No significant intra-day variation was seen, and it was excluded that the flux was modulated significantly by interstellar scintillation (Bremer et al. 1998). In the case of GRB 971214, no source was detected at 850 μm at the James Clerk Maxwell Telescope. The upper limit was 1 mJy on Dec 17-22 (Smith and Tilanus 1998).

3.6. Detection of a radio counterpart

After two unsuccessful searches for transient radio emission in GRB 970111 and GRB 970228 (Frail et al. 1997a), prompt VLA observations of the GRB 970508 error box allowed detection of a variable radio source within the error box, at 1.4, 4.8 and 8.4 GHz. It coincided with the optical transient and displayed a bizarre behaviour, in the sense that it remained at a steady level of brightness, flaring occasionally by a factor of two or three (Frail et al. 1997b), supporting the idea of the host galaxy as an active galactic nucleus. VLBI observations did not resolve the source (Taylor et al. 1997). Frail et al. proposed that the fluctuations could be the result of strong scattering by the irregularities in the ionized Galactic interstellar gas. The damping of the fluctuations with time indicates that the source expanded to a significantly larger size. The source was also detected at 15 GHz (Pooley and Green 1997).

All these observational efforts have provided the complete picture of a GRB multiwavelength spectrum, in the case of GRB 970508 (Fig. 5).

The observational characteristics of the three GRB counterparts (or *hypernovae*, a name proposed by Paczynski) can be accommodated in the framework of the fireball models (Mészáros et al. 1994, Mészáros and Rees 1997, Tavani 1997, Vietri 1997, Waxman 1997, Wijers et al. 1997), in which a compact source




Figure 5. The multiwaveband spectrum of OT/GRB 970508 on ~ 20 May 1997. Based on observations obtained at the VLA, Ryle, IRAM PdBI, ISO, Keck and 6-m SAO (Frail et al. 1997, Pooley and Green 1997, Bremer et al. 1998, Hanlon et al. 1998, Morris et al. 1997 and Sokolov et al. 1998).

releases 10^{53} ergs of energy within dozens of seconds. The plasma accelerates to relativistic velocities (the fireball). The blast wave is moving ahead of the fireball, and sweeps up the interstellar matter, producing an afterglow at frequencies gradually declining from X-rays to visible and radio wavelengths. Although some of the predicted behaviour has been observed, there are still many open questions (see Katz and Piran 1998).

4. SUMMARY

The existence of an X-ray afterglow in *all* bursts seems to be confirmed. The first optical/IR counterparts have been found in 1997. GRB 970508 was also detected in radio and mm, but prompt searches at other wavelengths failed to detect GRB 970111, GRB 970402 and GRB 970828.

BeppoSAX and *RossixTE* have opened a new window in the GRB field and it is widely accepted now that most GRBs, if not all, lie at cosmological distances. It is expected that *BeppoSAX*, *RossixTE* and *CGRO* will facilitate the discoveries of other counterparts and, together with the new high-energy observatories (*AXAF*, *SPECTRUM X/T*, *XMM*, *INTEGRAL*, *HETE 2*) and other satellites of the future *4th Interplanetary Network*, will definitively solve the long-standing Gamma-Ray Burst mystery.

5. ACKNOWLEDGEMENTS

I am grateful to many colleagues for very fruitful discussions, in particular to S. Brandt, M. Cerviño, E. Costa, M. Feroci, G. Fishman, F. Frontera, S. Golenetskii, J. Greiner, J. Gorosabel, S. Guzyi, R. Hudec, K. Hurley, J. Isern, C. Kouveliotou, N. Lund,

L. Metcalfe, H. Pedersen, M. de Santos, A. Shlyapnikov, R. Sunyaev, W. Wenzel, B. Wilson, C. Wolf, W. Wenzel, A. Yoshida and many others. I should also mention here the continuous support of my wife, M. E. Alcoholado-Feltström, who always patiently awaits for my return when sometimes I run off to the nearest observatory in order to perform the GRB follow-up observations and further 40-hr non-sleeping data analysis.

REFERENCES

- Aglietta, M. et al. 1995, *A&SS. Sci.* 231, 327
 Ali, S., Schaefer, R. K., Limon, M. and Piccirillo, L. 1997, *ApJ* 487, 114
 Antonelli, L. al. 1997, *IAU Circ.* 6792
 Atteia, J.-L., Barat, C., Hurley, K. et al. 1985, *A&A* 152, 174
 Atteia, J.-L. et al. 1987, *ApJS* 64, 305
 Balsano, R. J. et al. 1996, in *Gamma-Ray Bursts*, eds. C. Kouveliotou, M. F. Briggs and G. J. Fishman, *AIP* 384, 575
 Barthelmy, S. D., Palmer, D. M. and Schaefer, B. E. 1994, in *Gamma-ray Bursts*, eds. G. J. Fishman, J. Brainerd and K. Hurley, *AIP* 307, 392
 Becker-Szendy, R. et al. 1995, *ApJ* 444, 415
 Bhat, P. N. et al. 1996, in *Gamma-Ray Bursts*, eds. C. Kouveliotou, M. F. Briggs and G. J. Fishman, *AIP* 384, 585
 Blaes, O., Todd, H., Antonucci, R., Hurley, K. and Smette, A. 1997, *ApJ* 479, 868
 Boër, M., Atteia, J.-L., Gottardi, M. et al. 1988, *A&A* 202, 117
 Boër, M. et al. 1997a, *ApJ* 481, L39
 Boër, M. et al. 1997b, *IAU Circ.* 6795
 Bond, H. 1997, *IAU Circ.* 6655
 Bonnell, J. T. and Klebesadel, R. W. 1996, in *Gamma-Ray Bursts*, eds. C. Kouveliotou, M. F. Briggs and G. J. Fishman, *AIP* 384, 977
 Bontekoe, T. R. et al. 1995 *Ap&SS* 231, 285
 Borovička, J., Hudec, R., & Deődoch, A., 1992, *A&A* 258, 379.
 Bremer, M. et al. 1998, *A&A* 332, L13
 Castander, F. et al. 1997, *IAU Circ.* 6791
 Castro-Tirado, A. J., Brandt, S., Lund, N., Lapshov, I. Y., Terekhov, O. and Sunyaev, R. A. 1994a, in *Gamma-Ray Bursts*, eds. G. J. Fishman, J. J. Brainerd and K. Hurley, *AIP* 307, 17
 Castro-Tirado, A. J., Brandt, S., Lund, N. and Guzyi, S. 1994b, in *Gamma-Ray Bursts*, eds. G. J. Fishman, J. J. Brainerd and K. Hurley, *AIP* 307, 404
 Castro-Tirado, A. J., Gorosabel, J., Wolf, C. et al. 1997a, *IAU Circ.* 6657
 Castro-Tirado, A. J., Gorosabel, J., Heidt, J. et al. 1997b, *IAU Circ.* 6598
 Castro-Tirado, A. J., Gorosabel, J., Benitez, N., Wolf, C., Fockenbrock, R. et al. 1998a, *Sci* 279, 1011
 Castro-Tirado, A. J., Metcalfe, L. P., Laureijs, R., et al 1998b, *A&A* 330, 14

- Castro-Tirado, A. J., Gorosabel, J., Bowyer, S., Korpela, E. and Hurley, K. 1998c, in preparation
- Cataneese, M. et al. 1996, in *Gamma-Ray Bursts*, eds. C. Kouveliotou, M. F. Briggs and G. J. Fishman, AIP 384, 598
- Chevalier, C., Ilovaisky, S., Motch, C. et al. 1981, A&A 100, L1
- Chevalier, C. et al. 1997, IAU Circ. 6663
- Connaughton, V. et al. 1996, in *Gamma-Ray Bursts*, eds. C. Kouveliotou, M. F. Briggs and G. J. Fishman, AIP 384, 603
- Connaughton, V. et al. 1997, IAU Circ. 6683
- Connors, A. and McConnell, M. 1996, in *Gamma-Ray Bursts*, eds. C. Kouveliotou, M. F. Briggs and G. J. Fishman, AIP 384, 607
- Costa, E., Frontera, F., Heise, J. et al. 1997, Nat 387, 783
- DeMuth, D., Marshak, M. & Wagner, G. 1994, in *Gamma-Ray Bursts*, eds. G. J. Fishman, J. J. Brainerd and K. Hurley, AIP Conf. Proc. 307, 475
- Diercks, A. et al. 1997, IAU Circ. 6791
- Djorgovski, S. G. et al. 1997, Nat 387, 876
- Drinkwater, M. J. et al. 1997, IAU Circ. 6600
- Fenimore, E. et al. 1993, Nat 366, 40
- Feroci, M., Antonelli, L., Guainazzi, M. et al. 1998, A&A (submitted)
- Fishman, G. J. and Meegan, C. A. 1995, ARA&A 33, 415
- Frail, D., Kulkarni, S., Hurley, K. et al. 1994, ApJ 437, L43
- Frail, D. et al. 1997a, ApJ 438, L91
- Frail, D., Kulkarni, Nicastro, L., Feroci, M. and Taylor, G. 1997b, Nat 389, 261
- Frontera, F., Costa, E., Piro, L. et al. 1998, ApJ 493, L67
- Fruchter, A., Livio, M., Macchetto, D. et al. 1997, IAU Circ. 6747
- Funk, B. et al. 1996, in *Gamma-Ray Bursts*, eds. C. Kouveliotou, M. F. Briggs and G. J. Fishman, AIP 384, 612
- Galama, T. et al. 1997, Nat 387, 479
- Galama, T. et al. 1998, in preparation
- Garcia, M. et al. 1997, IAU Circ 6661
- Gorosabel, J. et al. 1998, (in preparation)
- Green, D. A. et al. 1996, in *Gamma-Ray Bursts*, eds. C. Kouveliotou, M. F. Briggs and G. J. Fishman, AIP 384, 617
- Greiner, J. and Motch, C. 1995, A&A 294, 177
- Greiner, J., Boër, M., Kahabka, P., Motch, C., Voges, W. 1995, in *Lives of Neutron stars*, eds. M. Alpar et al., NATO ASI C450, Kluwer, 519
- Greiner, J. 1996a, in *Multifrequency behaviour of high energy cosmic sources*, eds. F. Giovanelly and L. Sabau-Graziati, Mem. Soc. Astron. It. 67, 429
- Greiner, J., Bade, N., Hurley, K., Kippen, R. M. and Laros, J. 1996b, in *Gamma-Ray Bursts*, eds. C. Kouveliotou, M. F. Briggs and G. J. Fishman, AIP 384, 627
- Greiner, J., Wenzel, W., Hudec, R., Spurny, P., Florián, J. et al. 1996c, in *Gamma-Ray Bursts*, eds. C. Kouveliotou, M. F. Briggs and G. J. Fishman, AIP Conf. Proc. 384, 622
- Greiner, J., Schwarz, R., Englhauser, J., Groot, P. J. and Galama, T. 1997, IAU Circ. 6657
- Grindlay, J. E., Wright, E. L. and McCrowsky, R. E. 1974, ApJ 192, L113
- Grindlay, J. E., Cline, T. L., Desai, U.D. et al. 1982, Nat 300, 730
- Groot P.J. et al. 1997, IAU Circ. 6584
- Groot P.J., Galama T.J., van Paradijs J. et al. 1998, ApJ 493, L27
- Guarnieri, A. et al. 1997, A&A 328, L13
- Halpern, J. et al. 1997, IAU Circ. 6788
- Halzen, F. et al. 1991, Nat 353, 807
- Hanlon, L. et al. 1998, in preparation
- Harrison, T. E., McNamara, B. J. and Klemola, A. R. 1996, AJ 107, 254
- Hartmann, D. H. 1995, *Flares and Flashes*, eds. J. Greiner, H. W. Duerbeck and R. E. Gershberg, LNP 454, 367
- Hjellming, R. M. and Ewald, S. P. 1981, ApJ 246, L137
- Hudec R. et al. 1987, A&A 75, 71
- Hudec R. 1993, *Astrophys. Lett. Comm.* 28, 359
- Hudec, R., Dědoch, A., Pravec, P., & Borovička J. 1994a, A&A 284, 839
- Hudec R. 1995, Ap&SS 231, 239
- Hurley, K., Dingus, B., Mukherjee, R. et al. 1994, Nat 372, 652
- Hurley, K., Li, P., Laros, J. et al. 1995, ApJ 445, 348
- Hurley, K., Li, P., Murakami, T. et al. 1996, ApJ 469, L105
- Hurley, K. 1998, in *Gamma-Ray Bursts*, eds. C. A. Meegan, R. Preece & T. Koshut, AIP (in press)
- Katoh, M. et al. 1984, in *High Energy Transients in Astrophysics*, ed. S. Woosley, AIP 115
- Katz, J. I. and Piran, T. 1998, in *Gamma-Ray Bursts*, eds. C. A. Meegan, R. Preece & T. Koshut, AIP (in press)
- Klebesadel, R., Olsen, R. and Strong, I. 1973, ApJ 182, L85
- Klose, S., Eisloffel, J. and Richter, S. 1996, ApJ 470, L93
- Klose, S. & Tuffs, R. 1997, IAU Circ. 6611
- Koranyi, D. M. et al. 1994, MNRAS 271, 51
- Krawczynski, H. et al. 1996, in *Gamma-Ray Bursts*, eds. C. Kouveliotou, M. F. Briggs & G. J. Fishman, AIP 384, 656
- Larson, S., McLean, I. and Becklin, E. 1996, ApJ 460, L95
- Lee, B., Akerlof, C., Ables, E. et al. 1996, in *Gamma-Ray Bursts*, eds. C. Kouveliotou, M. F. Briggs & G. J. Fishman, AIP 384, 671
- LoSecco, J. 1994, ApJ 425, 217
- Mannheim, K., Jartmann, D. and Funk, B. 1996, ApJ 467, 532
- Marshall, F. et al. 1997a, IAU Circ. 6683
- Marshall, F., Cannizzo, J. K. and Corbet, R. H. D. 1997b, IAU Circ. 6727
- Masetti, N. et al. 1998, in *Gamma-Ray Bursts*, eds. C. A. Meegan, R. Preece and T. Koshut, AIP (in press).

- Mazets, E. P. et al. 1981, *Ap&SS* 80, 3
- McNamara, B. and Harrison, T. E. 1994, *AJ* 107, 1825
- Meegan, C. A., Fishman, G. J., Wilson, R. et al. 1992, *Nat* 355, 143
- Mészáros, P. and Rees, M. J. 1993, *ApJ* 418, L59
- Mészáros, P., Rees, M. J. and Papathanassiou, H. 1994, *ApJ* 432, 181
- Mészáros, P. and Rees, M. J. 1997, *ApJ* 476, 232
- Metzger, M. R. et al. 1997a, *IAU Circ.* 6655
- Metzger, M. R. et al. 1997b, *Nat* 387, 878
- Mignoli, M. et al. 1997, *IAU Circ.* 6661
- Morris, M. et al. 1997, *IAU Circ.* 6666
- Murakami, T. et al. 1991, *Nat* 350, 592
- Murakami, T. et al. 1996, *PASJ* 48, L9
- Murakami, T., Ueda, Y., Yoshida, A. et al. 1996, *IAU Circ.* 6732
- Murakami, T. et al. 1998, in *Gamma-Ray Bursts*, eds. C. A. Meegan, R. Preece & T. Koshut, *AIP* (in press)
- Nemiroff, R. 1994, *Comments Astrophys.* 17, 189
- Paczynski, B. and Xu, G. 1994, *ApJ* 427, 708
- Paczynski, B. 1998, *ApJ* 494, L45
- Padilla, L. et al. 1997, *Proc. XXV ICRC* 3, 57
- Palmer, D. et al. 1995, *A&SS* 231, 315
- Pedersen, H., Motch, C., Tarengi, M. et al. 1983, *ApJ* 270, L43
- Pedersen, H. et al. 1998a, *ApJ* 496, 311
- Pedersen, H. et al. 1998b, *A&A* (in preparation)
- Pedichini, F., Di Paola, A., Stella, L. et al. 1998, *A&A* 327, L36
- Pian, E. et al. 1998, *ApJ* 492, L103
- Piro, L. et al. 1997a, *IAU Circ.* 6617
- Piro, L. et al. 1997b, *IAU Circ.* 6656
- Pizzichini, G., Gottardi, M., Atteia, J.-L. et al. 1986, *ApJ* 301, 641
- Plunkett, S. et al. 1995, *A&SS* 231, 271
- Pooley, G. and Green, D. 1997, *IAU Circ.* 6670
- Remillard, R., Wood, A., Smith, D. and Levine, A. 1997 *IAU Circ.* 6726
- Sahu, K., Livio, M., Petro, L. et al. 1997, *Nat* 387, 476
- Schaefer, B. E. et al. 1984, *ApJ* 286, L1
- Schaefer, B. E., Cline, T. L., Desai, U. et al. 1987, *ApJ* 313, 226
- Schaefer, B. E., Cline, T. L., Atteia, J.-L. et al. 1989, *ApJ* 340, 455
- Schaefer, B. E. 1994, in *Gamma-ray Bursts*, eds. G. J. Fishman, J. J. Brainerd & K. Hurley, *AIP* 307, 382
- Schaefer, B. E. et al. 1994, *ApJ* 422, L71
- Schaefer, B. E., Cline, T. L., Hurley, K. and Laros, J. 1997a, *ApJ* 489, 693
- Schaefer, B. E. et al. 1997b, *IAU Circ.* 6658
- Smith, D. A. et al. 1997, *IAU Circ.* 6718
- Smith, I. A. and Tilanus, R. P. J. 1998, in preparation.
- Sofitta, P. et al. 1997, *IAUC* 6797
- Soifer, B. et al. 1997, *IAU Circ.* 6619
- Sokolov, V. V. et al. 1996, *Astron. Lett.* 22, 563
- Sokolov, V. V., Kopylov, A. I., Zharikov, S. V., et al. 1998, in *Gamma-Ray Bursts*, eds. C. A. Meegan, R. Preece & T. Koshut, *AIP* (in press)
- Stacy, J. G. et al. 1996, in *Gamma-Ray Bursts*, eds. C. Kouveliotou, M. F. Briggs and G. J. Fishman, *AIP* 384, 702
- Stecker, F. W. and de Jager, O. C. 1997, *ApJ* 476, 712
- Štěpán, P. & Hudec, R. 1996, *A&A* 305, 869
- Tavani, M. 1997, *ApJ* 483, L87
- Tanvir, N. et al. 1997, *IAU Circ.* 6796
- Taylor, G. B. et al. 1997, *Nat* 389, 263
- Trombka, J. I., Eller, E. L., Schmaderbeck, R. L., Adler, I., Metzger, A. E. et al. 1974, *ApJ* 194, L27
- van Paradijs, J. et al. 1997, *Nat* 386, 686
- Vietri, M. 1997, *ApJ* 488, L105
- Vrba, F., Hartmann, D. and Jennings, M. 1995, *ApJ* 446, 115
- Vrba, F. 1996, in *Gamma-Ray Bursts*, eds. C. Kouveliotou, M. F. Briggs and G. J. Fishman, *AIP* 384, 565
- Waxman, E. 1997, *ApJ* 489, L33
- Wdowczyk, J., Tkaczyk, W. and Wolfendale, A. W. 1972, *J. Phys. A*, 5, 1419
- Wheaton, W. A. et al. 1973, *ApJ* 185, L57
- Wijers, R. A. M. J., Rees, M. J. and Mészáros, P. 1997, *MNRAS* 288, L51
- Yoshida, A. and Murakami, T. 1994, in *Gamma-Ray Bursts*, eds. G. J. Fishman, J. J. Brainerd and K. Hurley, *AIP* 307, 333